INSECTICIDES INFORMATION

Elite Pest Management Ltd

To the Client:

You are advised that the product is approved for the use in domestic/commercial premises under the Control of Pesticides Regulations 1986. Your co-operation is requested in taking the following precautions to further minimise the risks to all occupants:

- **Rooms sprayed with a FICAM based insecticide can only be reoccupied by children or animals once the sprayed surfaces are dry**
- **Contact the Pest Control Officer for advice or assistance on the above telephone number**
- **In the case of accidental poisoning of animals, consult a veterinary surgeon taking this form with you**
- In case of any human accidental contact, seek medical advice taking this form with you.

The product marked on the list below has been used on your premises by our Pest Control Technician.

Insecticides Used (✓)

Product	Active Ingredient	
Ficam D	Bendiocarb	☐
Ficam W	Bendiocarb	☐
Ficam Plus	Bendiocarb	☐
Crackdown	Deltamethrin	☐
K-Othrine Rapide	Deltamethrin	☐
Other	Please Specify	☐

.....

These are anticholinesterase compounds.

Advice Helplines

Your doctor can obtain further advice from:

The Newcastle Poisons Information Bureau – Tel (0191) 2820300

Your veterinary surgeon can obtain further advice from:

Veterinary Poisons Information Service (VPIS) Tel: 020 71880200

All pesticides are poisonous; keep children, animals away from them.

HAZARDS AND EFFECTS:

DO NOT BREATH DUST OR SWALLOW

Inhalation & Swallowing	May cause dizziness, headache, excessive sweating, abnormal pains, tearfulness, slow heartbeat, muscular twitching, convulsions
Skin Contact	Can absorbed through the skin
Eye Contact	May cause constriction of the pupils, leading to temporary impaired vision

FIRST AID

Inhalation	Remove to fresh air, support if necessary, seek medical advice.
Swallowing	Wash mouth with water. Keep patient at rest, induce vomiting ONLY if patient is FULLY CONSCIOUS. Seek medical advice.
Skin	Remove contaminated clothing and wash affected area with soap and water.
Eyes	Rinse immediately with water for at least 15 minutes and obtain medical advice

NOTE TO PHYSICIAN

Contains anticholinesterase cabamates. Recommended antidote is atropine sulphate 2mg adult dose, preferably by injection, repeated as necessary until patient is fully atropinised. Do not use pralidoxime. All cases should be treated as an emergency. Respiratory failures the usual cause of death. If respiratory depression occurs, ventilation must be maintained.

NOTE TO OCCUPIERS

Ficam W treated rooms can only be reoccupied by children, pets, and other animals once the sprayed surfaces are dry.

FOR FURTHER ADVICE CONTACT:

Elite Pest Management Ltd
Elite House, Somerby Way
Heapham Road Industrial Estate
Gainsborough
Lincs
DN21 1QT
Tel: 01427 613923 or Mick Kilburn: 07957 485071 or Wayne Dunbar: 07947 114351